Google Custom Search

RSS feeds | Newsletter

GO

ART & STYLE ENTERTAINMENT COMPENDIUM

Physics | Chemistry | Geology and palaeontology | Biology | Environment | Astronomy | Health | Technology | — In pictures | — InMotion

Today's themes: cancer · brain · galaxy · rhino

Where am I? > Home > News > Health

Tags: care, diagnosis, health, hospital, illness, nurse, patient, physician

Professionals think that there aren't criteria for certifying that an illness is terminal

Science Centric | 14 July 2010 11:43 GMT

Brugada Syndrome test

Affordable 5-gene test for inherited Brugada syndrome genedx.com/cardiology

Ask a Doctor Online Now 9 Doctors Are Online. Ask a Question, Get an Answer ASAP

Open Days
Take the initiative University of Plymouth

Online University Degree
Find College & University Programs by Type of Degree.

Ads by Google

Physicians that have to decide whether an illness is terminal think that 'there are not valid and adequate criteria for certifying that an illness is terminal. This is the conclusion drawn from a pioneer study conducted at the University of Granada, and recently published in the International Journal of Clinical and Health Psychology (IJCHP).

For the purpose of this study, 42 health professionals were thoroughly interviewed. Of this group, 21 were physicians and 21 were nurses working with terminally-ill patients in hospitals in the province of Granada. Of the 42 participants, 22 were women and 20 were men with ages between 23 and 52. Of the participants, 17 worked in community health centres, 18 worked in public hospitals, 4 in combined units and 3 worked in private health centres

This study revealed interesting findings. The answers given by the professionals interviewed revealed that the diagnosing of terminal illness in their daily work differs depending on the type of health centre - primary or specialised care - and the type of professional - physician or nurse.

Concretely, those professionals working in community health centres - both physicians and nurses - use and record 'terminal illness' diagnosis to define the clinical condition of their patients. On the other hand, the answers provided by professionals working in public hospitals differ significantly: while nurses use euphemisms or synonyms rather than 'terminal illness' diagnosis, physicians use this diagnosis in professional circles and, although they are aware of this circumstance, they avoid using it in their medical reports

This research conducted at the University of Granada reveals that health professionals have difficulties in certifying terminal illnesses. The question emerges on whether such situation could be preventing palliative care from being provided to a significant number of terminal patients. This means that many terminal patients are being deprived from professional assistance aimed at helping them in facing the end of their lives, and at providing them an acceptable quality of life. Additionally, professionals have a distorted view of the role and purpose of palliative assistance, since they are emotionally conditioned by a diagnosis that - in our society - is associated to a 'death

This study was co-authored by Maria Paz Garcia Caro, Francisco Cruz Quintana, Jacqueline Schmidt Rio Valle, Antonio Munoz Vinuesa, Rafael Montoya Juarez, Diego Prados Pena y Miguel C. Botella Lopez from the University of Granada, and Atthanasios Pappous, from the University of Kent, United Kingdom.

For a deeper understanding of the difficulties that a diagnosis of 'terminal illness' involves and its consequences, health professionals were asked what illnesses get diagnosed as 'terminal' and when such diagnosis is disclosed. For most physicians and nurses, 'terminal illness' diagnosis is associated only to patients with cancer. However, a significant number of professionals working in community health centres also associate this diagnosis to patients with an advanced stage of some chronic and/or degenerative non-cancerous illness.

As regards terminal diagnosis, it is mostly associated to patients in preagony or agony phase. This opinion is shared by most professionals working both in hospitals and in community health centres.

The study emphasises the fact that, although the criteria and references for diagnosing a terminal illness - at least in oncological cases - were defined 20 years ago, such diagnosis is still employed in a very restrictive way. Diagnosis of a terminal illness involves a change of treatment, and when patients in terminal stage are not diagnosed as such, they and their families

BOILING POINT

 McDonald's recalls Shrek glasses due to potential — <u>incolorated steals Silter glasses due to potential cadmium risk</u> — The U.S. Consumer Product Safety Commission (CPSC) just announced a recall of 'Shrek Forever After 3D' Collectable Drinking...

— <u>Hogchoker - the new Internet star</u> — A small flatfish living along the coast of North America is the new Internet star.

Currently the hotness for this particular...

— Cancer deaths are projected to double by 2030 — Cancer deaths are projected to double in the next two decades. A report issued by the International Agency for Research on...

More Boiling point ..

MINUSCULE

— 'Brain recycling' puts kids' writing in a twist — When children learn to read and write, they often get things...

— Camel spiders are sticky killers — Camel spiders have evolved a unique way to capture their insect...

— Can a gene test tell you whether you'll live to 100? — If a genome test could predict your odds of living to 100, would...


— Super squid says organ discovered — The mating habits of decrease squid have been revealed for the deep-sea squid have been revealed for the...

More Minuscule..

RSS feeds, newsletter

Find the topic you want. Science Centric offers several RSS feeds for the News section.

Or subscribe for our Newsletter, a free e-mail publication. It is published practically every day.



New method may accelerate drug discovery for difficult diseases like Parkinson's -Whitehead Institute scienti have developed a rapid, inexpensive drug-screening

method that could be used to



Caffeine reverses memory another reason to pour that extra cup. When aged mice bred to develop symptoms of Alzheimer's disease...



Site for alcohol's action in the brain discovered — Alcohol's inebriating effects are familiar to everyone. But the molecular details of alcohol's impact on brain activity remain



The battle for CRTC2: How obesity increases the risk for diabetes — Obesity is probably the most important factor in the development of insulin resistance, but science's understanding of the...

are deprived from the special care and assistance they need

Similarly, the study remarks the lack of studies on the emotional charge suffered by health professionals treating terminal patients. This study states that this is a key factor when explaining the restrictive use of terminal illness diagnosis, especially in hospitals

The research conducted at the University of Granada emphasises the need to improve professional training of care providers in diagnosing a terminal illness, especially in chronic and degenerative illnesses in advanced stages. Additionally, the study remarks the need of further research for determining the scientific parameters on which terminal diagnosis must be based.

Source: Universidad de Granada

Nowicky Pharma • Ukrain
Alternative treatment New hope for patients

Brugada Syndrome test

Affordable 5-gene test for inherited Brugada syndrome

New AUA Campus

Caribbean Medical school opens most modern campus in the region.

Ads by Google

Leave a comment

The details you provide on this page [e-mail address] will not be used to send unsolicited e-mail, and will not be supplied to a third party! Please note that we can not promise to give everyone a response. Comments are fully moderated. Once approved they will be posted within 24 hours.

| , . | | |
|--------------------|---|--|
| Your name: | | |
| | | |
| Your e-mail: | | |
| | | |
| Comment: | | |
| Comment | Ī | |
| | | |
| | | |
| | | |
| | | |
| | | |
| | | |
| 1753 | | |
| Verification code: | | |
| Volumbation code: | | |
| (| | |
| SEND | | |

More on Science Centric News | Health

Cancer: The cost of being smarter than chimps?

— Are the cognitively superior brains of humans, in part, responsible for our higher rates of cancer? That's a question that has nagged at John McDonald, chair of Georgia Tech's School..

Study gives clues to how adrenal cancer forms

— At the ends of chromosome are special pieces of DNA called telomeres. Think of it as the little tip that caps off a shoelace. The telomeres send signals to the cells to let them know...

GO





LATEST HEADLINES

Prediction tool helps estimate local recurrence in patients with noninvasive breast cancer Many physicians do not accept responsibility

to report incompetent, impaired colleagues New treatment for crippling diabetic Charcot

Lessons from efforts to reduce hospitalacquired infections Medications found to cause long term

cognitive impairment of aging brain Help is on the phone: Reducing pain and depression of cancer

More Health...

Popular tags in Health: cancer · diabetes · malaria · obesity

Google Custom Search

Science Centric — Science, health and technology, breaking news

Front page | News | Travel | Art & style | Entertainment | Compendium | Resources | Site map | Accessibility | About us | Contact us | RSS feeds

Copyright © 2007-2010 Science Centric.

Science Centric delivers breaking news about the latest scientific discoveries in the fields of physics, chemistry, geology and palaeontology, biology, environment, astronomy, health, and technology. Use of this website assumes acceptance of Terms of use and Privacy policy.

Lepidopterology | National Museum of Natural History, Sofia