


[Health News](#)
[Videos](#)
[Article Opinions](#)
[Advertise](#)
[Contact](#)

 Personalization
[login](#) | [register](#)


Search

 powered by

Follow us on:



Pharma Industry / Biotech Industry News

[Useful Links](#)

Drug Encapsulation And Gene Therapies Use New Data Obtained On Liposomes

 Main Category: [Pharma Industry / Biotech Industry](#)

 Also Included In: [Genetics](#); [IT / Internet / E-mail](#)

Article Date: 09 Feb 2011 - 3:00 PST

3

[email to a friend](#) [printer friendly](#) [opinions](#)

University of Granada scientists and the Spanish Higher Institute for Scientific Research (CSIC) have made significant progress in understanding lipid membranes, which are extensively employed in the development of cosmetic and drug products, and which have potential application in the field of nanotechnology.

Phospholipid vesicles (liposomes) are colloidal systems that arise considerable interest from the pharmaceutical, cosmetic and food industry, since they are biocompatible in protein, nucleic acid, drug, etc encapsulation. Further, from a scientific perspective, liposomes are considered a model system of cell membranes that have been implemented in the study of biological transport processes through cell membranes, as well as in the study of aggregation processes induced by biological substances, etc.

Current Article Ratings:

Patient / Public:	Not yet rated
Healthcare Prof:	Not yet rated

 Find other articles on: ["University of Granada"](#)

To develop products of biotechnological interest, understanding thoroughly the electrostatic properties of these membranes is necessary. This was the purpose of Alberto Martín Molina and César Rodríguez Beas, from the Department of Applied Physics of the University of Granada, and Jordi Faraudo from the Instituto de Ciencias de Materiales in Barcelona (CSIC), authors of a study recently published in the journal *Physical Review Letters* (Vol. 104; pp 168103 (2010). 104, pp 168103 (2010)).

Inverting its Electrostatic Charge

This study discloses why certain lipid membranes can invert their surface electrostatic charge, that is, why these membranes have negative charge, but they can function as positive charge material in specific circumstances. This type of membranes are extensively employed in gene therapies.

Such behaviour is due to the fact that the interphase of these membranes in water is soft, permeable and highly hydrated. "Such environment attracts small objects with significant electric charge. These membranes tend to gather in large groups acquiring electric charge", researchers state. For the purpose of this study, electrophoresis experiments and computer-based simulations were conducted. Such trials were made using a supercomputer belonging to the Spanish Supercomputing Centre, since these trials required a long time and high calculation performance.

After a sustained period of several months, researchers obtained revealing results from simulations, which allowed them to prepare a new inversion mechanism for their experimental system. This mechanism is as follows: phospholipid membranes have the ability to absorb lanthanum cations, which go from being associated to the solution water molecules to associating to the membrane atoms.

Source:
 Alberto Martín Molina
 University of Granada

 Please rate this article: [Patient / Public:](#) or [Health Professional:](#)

(Hover over the stars then click to rate)

Ads by Google

[Life Sciences QMS](#) - Lean Quality Management Systems Ensuring Regulatory Compliance - [www.gxpi.com](#)