Truth About Drugs Free Truth About Drugs Info Kit & Booklets. Learn The Facts Now! DrugFreeWorld.org Psychiatric Drug Facts Do You Know The Long Term Effects Of Psychiatric Drugs? Free Info Kit CCHR.0

ADVERTISEMENT

24.com Home | Blogs | Albums | Classifieds | **24.com sites** ▼

Are you a medical whizz?

Check your medical knowledge right here with this guiz.

Lunchtime workout

A lunch-time quickie might be all you have time for

Subscribe to our newsletters | Win!

- Hampers worth R2000
- R 1000 a month









Health News Conditions A-Z Your Body Focus Lifestyle Diet & Nutrition Fitness Sex Experts Community Tests & Tools Medical Schemes Win

Weekly Poll

Free-range, organic, grass-fed: do you know what you're eating?

- O Yes, I keep up with research
- O Sort of, I read labels
- l'd like to, but I don't always know the difference
- No I eat what I like
- O I don't know about this

SUBMIT

General **Newsletter**

Sign up for our newsletter to get all the latest Health

email address

YOU ARE IN > News > Addiction & substance abuse

6 in 10 drug addicts abuse partner

Last updated: Friday, March 04, 2011

Be the first of your Recommend friends to recommend

About six out of 10 male drug-abusers direct some type of violence against their intimate partners, according to a study conducted at the Department of Pedagogy of the University of Granada. The study revealed a high rate of domestic violence - both

The study also detailed the most recurrent forms of abuse, as well as the variables associated to them

pysical and psychological - by male drug-abusers against women.

The study revealed that between 6.5% and 72.4% of the population admits to having committed some form of violence against their partner. The less prevalent conduct was "I forced her to abort against her will" (6.5%), and the most prevailing being "I show total disregard for her needs, wishes and interests".

In addition, 63.5% of the respondants ensured that they wanted to know what their partner makes at all times of the day, their hours, whom they speak to". The rate of physical violence oscillates between 6.5% and 21%, while psychological violence varies between 7.3% and 72.4%.

Aware of voilence

The study revealed that 51% of male drug-abusers are aware of their committing violence against their partners and, although they know that such violence has a deep impact on their partners, they are not willing to break off, or to let their spouses abandon them, thus resorting to whatever strategies are necessary

Additionaly, psychological violence is more frequent than physicial violence in this collective. The most representative forms of violence being, by order of incidence: personal control, sexual abuse, emotional neglect, emotional blackmail out of guilt, disregard for their ideology or religion, abuse based on gender roles, emotional blackmail through enforcement, economic abuse, emotional blackmail out of fear, loss of personal control and social isolation.

This research was carried out by Amelia Matute López and professor Andrés Soriano Díaz. The author took a sample of 153 men aged between 20 and 65 years, attending some of rehabilitation units in Andalusia; all of them were cocaine, alcohol, heroin or heroin-cocaine abusers.

The study was a questionnaire-based prevalence survey of spousal abuse, containing also some questions to obtain some information on the variables selected.

78.8% of couples seperate, get back together

The research conducted at the University of Granada revealed that 78.8% of the couples ended in separation, although such separation was generally temporary. In most of the cases, the couple ended getting back together

"Despite their negative effect on their health and well-being, women remain in this type of relationship as a result of the pressure exerted by their male partners and of the romantic myths of unconditional and imbalanced love," Amelia Matute states.

This research contributes to better understanding the phenomenon of gender-based violence, in a sector of the population that had not been previously studied. In Spain, there are some studies based on the general population, but not on specific sectors of the population that could be considered of high-risk, such as it is the case of drug-abusers, as this study reveals. - (EurekAlert!, March 2011)

Today's top stories

Are men with deep voices cheaters?

Genital size linked to male fertility


Health dept faces malpractice suits

Happy hour linked to pub violence Unethical research retracted

Fear battles Aids fatalism

Drug use linked to plane crashes

Is Computer Vision Syndrome real?



Profmed

Blogs

People are blogging about

Clicks Pharmacy

Forums

Find out what others are

Your Body ADHD Acne Allergy Alzheimer's

1 de 2 07/03/2011 10:30