



Log in

¡FELICITACIONES! ¡Usted es el GANADOR del iPad 2!

Haga clic en el botón "Sí" a continuación para reclamarlo antes **de que se acabe el tiempo.**

Sí

No

HOME

VIDEO

NEWS

IMAGES

HEALTH

EDUCATION

TOPICS

MOBILE

Search

Space

Science

Technology

Health

General

Sci-Fi & Gaming

Oddities

International

Business

Education

Depression And Pain Increase Fatigue In Breast Cancer Survivors

September 14, 2011

Like



Twitter

0



Email

0



Share

0



Digg

0

The study has been published in the [European Journal of Cancer Care](#)

In Spain, 5-year survival following breast cancer diagnosis is more than 83%. Around 66% suffer fatigue following treatment. A Spanish research establishes the factors associated with tiredness in cancer survivors to improve their quality of life and rehabilitation.

"Cancer-related fatigue is the symptom that most limits quality of life and is most common in patients that survive cancerous processes," explained Manuel Arroyo, researcher from the University of Granada and lead author of a study that links psychological disorders and physical pain episodes with fatigue after treating a breast tumor.

More than 66% of breast cancer survivors suffer tiredness following recovery, caused directly by the disease, physical deterioration or the treatment received. Therefore, understanding the factors related with fatigue and how they can be alleviated optimizes survivors' recovery.

Fifty-nine female patients treated for breast cancer were followed-up one year after having clinically overcome the disease. The researchers assessed their psychological and physical condition as well as different aspects linked to the typical medical symptoms following a cancerous process (tiredness, pain, limited movement, depression, etc.).

A statistical procedure (resampling) allowed inferences to be made similar to those that would be obtained from larger samples. "This method means that the data were more reliable and eliminated the problem of having a reduced sample size," explained Arroyo. "It is difficult to find volunteers because patients are not often very willing to participate in research after having been through such harsh treatment."

The results show that the patients most affected by tiredness following treatment also suffer episodes of depression and body image deterioration, neck and shoulder pain, and limited arm movement, possibly due to the surgical intervention.

Effects of survival

Following breast cancer treatment, patients present with physical and psychological symptoms that influence their health.

Previous studies have already observed self-esteem- and body image-related disorders following the cancerous process. But for the first time, a team of researchers has associated sensory hypersensitivity, limited movement and certain psychological conditions with fatigue observed following cancer treatment.

"These findings should motivate patient support programs which improve their psychological condition and offer resources that can reduce pain," pointed out Arroyo, who further stressed that "if fatigue is not treated, patients can suffer it for years, having a serious physical, emotional, social and economic impact."

—

On the Net:

[FECYT – Spanish Foundation for Science and Technology](#)
[European Journal of Cancer Care](#)

Source:



Post a new comment

Login

Post

0 Comments

Related Articles

[Common Invasive Test Not Necessary For Kidney Disease Patients](#)

[New Strategy Likely To Speed Drug Development For Rare Cancers](#)

[Damaged Gait And Balance Can Recover With Long-Term Abstinence From Alcohol](#)

[People Born After World War II Are More Likely To Binge Drink, Develop Alcohol Disorders](#)

[Chronic Drinking Leads To Reduced Cortical Thickness In Frontal And Temporal Brain Regions](#)