

- [My news](#)
- [Unread news](#)
- [Science wire](#)

•

- Reading Phys.org on your mobile device? Enjoy a convenient format at your fingertips. Phys.org can now be found on Google Currents (Android and iOS). [Read more...](#)

# UGR and Michael J. Fox Foundation take a step forward in the study of Parkinson's disease

November 9th, 2012

During the past year, the research group TIC 218 at the Department of Signal Theory, Telematics and Communications attached to the University of Granada and the Andalusian Information Technology and Communications Center (CITIC) has implemented the methods for the detection and diagnosis of Alzheimer's disease, within the PPMI initiative of the Michael J. Fox foundation for the study of Parkinson's disease. In addition, other databases were used such as that of the Virgen de la Victoria Hospital, Málaga, Spain, and Virgen de las Nieves Hospital, Granada, Spain.

The research studies were financed by the Andalusian Regional Government through its excellence research program calls TIC-4530, TIC-2566 and TIC-7103. Additional funds were granted by the Spanish National R&D Plan call TEC2008-02113 and "DiagnoSIS" TEC2012-34306. The research group has been awarded several prizes as the University of Granada Premio del Consejo Social and the Andalusian Regional Government's Premio Andalucía Sociedad de la Información.

This research group has published more than 40 papers indexed in the ISI-JCR in the field of diagnosis and scan imaging analysis. The last three papers –devoted to the study of Parkinson's disease- have been published in the most prestigious journals in the field of neuroscience such as *Medical Physics* (impact index 3.075) and *NeuroImage* (impact index 5.985, N°1 journal in its category).

## Understanding the behavior of brain functions in patients with Parkinson's disease

The introduction of SPECT imaging worldwide for the diagnosis of Parkinson's disease (PD) –based on the innovative DaTSCAN agent– has allowed experts to better understand the behavior of behavior of brain functions in patients diagnosed with Parkinson's disease. The data provided by SPECT images can be processed by pattern-recognition machine, which allow an objective analysis of data.

## A collaborative work with Virgen de la Victoria Hospital

The latest study conducted by this research group (Automatic assistance to Parkinson's disease diagnosis in DaTSCAN SPECT imaging. I. A. Illán, J. M. Górriz, J. Ramírez, F. Segovia, J. M. Jiménez-Hoyuela, and S. J. Ortega Lozano *Med. Phys.* 39, 5971 (2012)) presents a computer procedure that does not need the intervention of an expert (normalization, location of relevant regions, "clinical eye"...). In addition, this new technique systematically analyzes the effect of the different elements involved in the building process of the computer-assisted diagnosis technique on the accuracy of the final diagnosis. As a result, the technique proposed may exceed 90% of accuracy in the diagnosis of PD with the right selection of elements, and with the additional advantage of being a robust and simple computer software.

**A Paper Accepted for the Prestigious Journal *Neuroimage***

In this study, stable-distribution parameters, concretely those of scale and position, are used to linearly convert the intensity values for each voxel. This conversion is performed in such a way that the new intensivity histograms for each brain image have the same values for the parameters position and dispersion. The proposed method has a better performance than the traditional intensity normalization method, which is based on the ratio between the intensity values of specific and non-specific brain regions.

Provided by University of Granada



CORPORACIÓN**CAPILAR**

HAIRCARE

## More news stories

## Neural circuit in the songbird brain that encodes representation of learned vocal sounds located