

HOME

Cloud Service Integration

www.BMC.com/Cloud

Debunking Myths About the Cloud Get the Straight Facts at BMC™

LIFESTYLE

HEALTH

SCIENCE



SPACE

WORK

Health Images

NATIONAL.

WHAT'S HOT:

BJP Congress Cricket Armstrong

WORLD

Exposure to pesticides in food, air and water ups diabetes risk

SPORTS

ENTERTAINMENT

Feb 6, 10:40 am



Washington, February 6(ANI): Scientists have found that there is a direct relationship between exposure to pesticides (Persistent Organic Pollutants, CPOs) in food, air and water and prevalence of type 2 diabetes in adults, regardless of age, gender and body mass index.

These substances tend to concentrate in body fat, and

they might be one of the reasons why obese people are more likely to develop diabetes, since the more fat the higher the COP concentrations in the body.

In a study conducted at the University of Granada, researchers demonstrate that people with higher concentrations of DDE -the main metabolite in the pesticide DDT- are four times more likely to develop type 2 diabetes than other people.

In addition, the risk of type 2 diabetes is also associated with exposure to beta-HCH (beta-Hexachlorocyclohexane), which is present in the formula of the pesticide Lindano.

To carry out this study, the researchers analyzed the concentrations of a specific group of COPs in the adipose tissue of 386 adult subjects assisted at San Cecilio hospital, Granada, and Santa Ana hospital, Motril, Spain.

According to the University of Granada and San Cecilio researcher, Juan Pedro Arrebola, "human adipose tissue (commonly known as "fat") acts as an energy reservoir and has an important metabolic function. However, adipose tissue can store potentially harmful substances, such as persistent organic pollutants (COPs)".

This makes COPs concentrations a useful marker of a subject's exposure to COPs. COPs are a group of chemicals with diverse characteristics, which are present in pesticides, industrial waste and building materials. These compounds penetrate the body mainly through food, but also through air or the skin.

ording to professor Arrebola, "the mechanism of action by which COPs increases the risk of diabetes is still unknown. However, some researchers have suggested that COPs might cause an unological response when they penetrate estrogen receptors in tissues associated with the abolism of sugars."

results were recently reported in the journal Environmental Research. (ANI)

Pesticides, type 2 diabetes, DDT, Persistent Organic Pollutants

10 Diabetes Secrets

TECH

Recommend 0 RealDiabetesTruth.com/Diabetes

Beat Diabetes Fast With These
10 Diabetes Secrets. Free
16pp Report



AdChoices [⊳

LATEST STORIES

1097787

- Why obese people are unable to lose weight?
- Vitamin D and omega-3 may help fight Alzheimer's
- Alcohol + diet drinks elevate intoxication more than alcohol + regular drinks
- Green tea and red wine may hold promise for Alzheimer's treatment
- Vitamin C supplements boost kidney stone risk in men
- Staying fit during middle age could ward off dementia
- Sunlight may help stave off rheumatoid arthritis in women
- Calcium supplements may be bad for a man's heart
- Low vitamin D levels may raise risk of Type 1 diabetes
- Diabetics treated with insulin 'at greater risk of health complications'



TOP VIDEO STORIES



Women don'... Stock Pulse: Fe...

Death risk from obesity increases with age Feb 6, 4:23 pm